

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 February 2002 (28.02.2002)

PCT

(10) International Publication Number
WO 02/15933 A2

- (51) International Patent Classification⁷: **A61K 45/00**
- (21) International Application Number: **PCT/EP01/09587**
- (22) International Filing Date: **20 August 2001 (20.08.2001)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
09/643,641 **22 August 2000 (22.08.2000)** **US**
- (71) Applicant (*for all designated States except AT, US*): **NOVARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).**
- (71) Applicant (*for AT only*): **NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H. [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).**
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **ALLISON, Malcolm [GB/CH]; Grenzacherstrasse 108, CH-4058 Basel (CH). GATLIN, Marjorie, Regan [US/US]; 913 Lovering Avenue, Wilmington, DE 19806 (US).**
- (74) Agent: **BECKER, Konrad; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).**
- (81) Designated States (*national*): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (*regional*): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
— *without international search report and to be republished upon receipt of that report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/15933 A2

(54) Title: **COMBINATION**

(57) Abstract: The present invention relates to a combination, especially a pharmaceutical composition, comprising as active ingredients (i) an AT₁-receptor antagonist or a pharmaceutically acceptable salt thereof; (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or (b) an insulin sensitizer or a pharmaceutically acceptable salt thereof; and, in case of a pharmaceutical composition, a pharmaceutically acceptable carrier.

BEST AVAILABLE COPY

Combination

The present invention relates to a combination, especially a pharmaceutical composition, comprising as active ingredients

- (i) an AT₁-receptor antagonist or a pharmaceutically acceptable salt thereof;
 - (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof; and,
- in case of a pharmaceutical composition, a pharmaceutically acceptable carrier.

AT₁-receptor antagonists (also called angiotensin II receptor antagonists or blockers) are understood to be those active ingredients that bind to the AT₁-receptor subtype of angiotensin II receptor but do not result in activation of the receptor. As a consequence of the blockade of the AT₁ receptor, these antagonists can, for example, be employed as antihypertensives or for treating congestive heart failure.

The class of AT₁ receptor antagonists comprises compounds having differing structural features, essentially preferred are the non-peptidic ones. For example, mention may be made of the compounds that are selected from the group consisting of valsartan (cf. EP 443983), losartan (cf. EP 253310), candesartan (cf. EP 459136), eprosartan (cf. EP403159), irbesartan (cf. EP 454511), olmesartan (cf. US 5616599), tasosartan (cf. EP 539086), and telmisartan (cf. EP 502314), or, in each case, a pharmaceutically acceptable salt thereof.


Preferred AT₁-receptor antagonist are those agents that have been marketed, most preferred is valsartan or a pharmaceutically acceptable salt thereof.

Insulin secretion enhancers are active ingredients that have the property to promote the secretion of insulin from pancreatic β -cells. Examples of insulin secretion enhancers are sulfonylureas (SU), especially those which promote the secretion of insulin from pancreatic β -cells by transmitting signals of insulin secretion via SU receptors in the cell membrane, including (but are not limited to) tolbutamide; chlorpropamide; tolazamide; acetohexamide; 4-chloro-N-[(1-pyrrolidinylamino)carbonyl]-benzensulfonamide (glycopyramide); glibenclamide (glyburide); gliclazide; 1-butyl-3-metanilylurea; carbutamide; glibonuride;

- 2 -

glipizide; gliquidone; glisoxepid; glybuthiazole; glibuzole; glyhexamide; glymidine; glypinamide; phenbutamide; and tolylcyclamide, or a pharmaceutically acceptable salt thereof.

Insulin secretion enhancers furthermore include short-acting insulin secretion enhancers, such as the new phenylalanine derivative nateglinide [N-(trans-4-isopropylcyclohexyl-carbonyl)-D-phenylalanine] (cf. EP 196222 and EP 526171) of the formula



repaglinide [(S)-2-ethoxy-4-{2-[[3-methyl-1-[2-(1-piperidiny)phenyl]butyl]amino]-2-oxoethyl}benzoic acid – cf. EP 589874] ; calcium (2S)-2-benzyl-3-(cis-hexahydro-2-isoinдолinylcarbonyl)-propionate dihydrate (mitiglinide – cf. EP 507534); furthermore representatives of the new generation of SUs such as glimepiride (cf. EP 31058); and in free or pharmaceutically acceptable salt form.

Insulin secretion enhancers likewise include the long-acting insulin secretion enhancer DPP-IV inhibitors, GLP1 and GLP1 agonists.

DPP-IV is responsible for inactivating GLP-1. More particularly, DPP-IV generates a GLP-1 receptor antagonist and thereby shortens the physiological response to GLP-1. GLP-1 is a major stimulator of pancreatic insulin secretion and has direct beneficial effects on glucose disposal.

The DPP-IV inhibitor can be peptidic or, preferably, non-peptidic. DPP-IV inhibitors are in each case generically and specifically disclosed e.g. in WO 98/19998, DE 196 16 486 A1, WO 00/34241 and WO 95/15309, in each case in particular in the compound claims and the final products of the working examples, the subject-matter of the final products, the pharmaceutical preparations and the claims are hereby incorporated into the present application by reference to these publications. Preferred are those compounds that are

specifically disclosed in Example 3 of WO 98/19998 and Example 1 of WO 00/34241, respectively.

GLP-1 is a insulinotropic peptide which was described, e.g., by W.E. Schmidt et al. in Diabetologia 28, 1985, 704-707 and in US 5,705,483.

The term "GLP-1 agonists" used herein means variants and analogs of GLP-1(7-36)NH₂ which are disclosed in particular in US 5,120,712, US 5,118,666, US 5,512,549, WO 91/11457 and by C. Orskov et al in J. Biol. Chem. 264 (1989) 12826. The term "GLP-1 agonists" comprises especially compounds like GLP-1(7-37), in which compound the carboxy-terminal amide functionality of Arg³⁶ is displaced with Gly at the 37th position of the GLP-1(7-36)NH₂ molecule and variants and analogs thereof including GLN⁹-GLP-1(7-37), D-GLN⁹-GLP-1(7-37), acetyl LYS⁹-GLP-1(7-37), LYS¹⁸-GLP-1(7-37) and, in particular, GLP-1(7-37)OH, VAL⁸-GLP-1(7-37), GLY⁸-GLP-1(7-37), THR⁸-GLP-1(7-37), MET⁸-GLP-1(7-37) and 4-imidazopropionyl-GLP-1. Special preference is also given to the GLP agonist analog exendin-4, described by Greig et al in Diabetologia 1999, 42, 45-50.

A preferred insulin secretion enhancer is repaglinide, most preferred is nateglinide.

The term nateglinide likewise comprises crystal modifications such as disclosed in EP 0526171 B1 or US 5,488,510, respectively, the subject matter of which, especially with respect to the identification, manufacture and characterization of crystal modifications, is herewith incorporated by reference to this application, especially the subject matter of claims 8 to 10 (being directed to the H-form crystal modification) as well as the corresponding references to the B-form crystal modification.

The structure of the active agents identified by generic or tradenames may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g. Patents International (e.g. IMS World Publications). The corresponding content thereof is hereby incorporated by reference. Any person skilled in the art is fully enabled to identify the active agents and, based on these references, likewise enabled to manufacture and test the pharmaceutical indications and properties in standard test models, both in vitro and in vivo.

- 4 -

The term "short-acting insulin secretion enhancer" comprises corresponding agents with a maximum secretion of insulin that is attained within one hour, preferably within 30 minutes, after the administration of the agent, most preferably within 20 minutes having a biological half-life, $T_{1/2}$, of less than two hours, preferably, 1.5 hours. The term long-acting insulin secretion enhancer" comprises corresponding agents with a maximum secretion of insulin that is attained more than one hour after administration of the agent.

A preferred insulin sensitizer is metformin or a pharmaceutically acceptable salt thereof such as the mono-hydrochloride.

Especially preferred is a combination of valsartan or a pharmaceutically acceptable salt thereof and nateglinide or a pharmaceutically acceptable salt thereof.

The corresponding active ingredients or a pharmaceutically acceptable salts thereof may also be used in form of a solvate, such as a hydrate or including other solvents, used for crystallization.

The compounds to be combined can be present as pharmaceutically acceptable salts. If these compounds have, for example, at least one basic center, they can form acid addition salts. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center. The compounds having an acid group (for example COOH) can also form salts with bases.

The pharmaceutical activities as effected by administration of representatives of the class of AT_1 -receptor antagonists or insulin secretion enhancers, respectively, or of the combination of active agents used according to the present invention can be demonstrated e.g. by using corresponding pharmacological models known in the pertinent art. The person skilled in the pertinent art is fully enabled to select a relevant animal test model to prove the hereinbefore and hereinafter indicated therapeutic indications and beneficial effects.

To evaluate the antihypertensive activity of the combination according to the invention, for example, the methodology as described by Lovenberg W: Animal models for hypertension research. Prog. Clin. Biol. Res. 1987, 229, 225-240 may be applied. For the evaluation that the combination according to the present invention may be used for the treatment of

congestive heart failure, for example, the methods as disclosed by Smith HJ, Nuttall A: Experimental models of heart failure. *Cardiovasc Res* 1985, 19, 181-186 may be applied. Molecular approaches such as transgenic methods are also described, for example by Luft et al.: Hypertension-induced end-organ damage. "A new transgenic approach for an old problem." *Hypertension* 1999, 33, 212-218.

The insulin secretion enhancing properties of the combination according to the present invention may be determined by following the methodology as disclosed, for example, in the publication of T.Ikenoue et al. *Biol.Pharm.Bull.* 29(4), 354-359 (1997).

The corresponding subject matter of these four references is herewith incorporated by reference in this specification.

Accordingly, the combination according to the present invention may be used, e.g., for the prevention, delay of progression or treatment of diseases and disorders that may be inhibited by the inhibition of AT₁ receptor, that may be inhibited by the enhancement of insulin secretion and that may be inhibited by insulin sensitization. Especially, the combination according to the present invention may be used, e.g., for the prevention, delay of progression or treatment of diseases and disorders selected from the group consisting of hypertension, congestive heart failure, diabetes, especially type 2 diabetes mellitus, diabetic retinopathy, macular degeneration, diabetic nephropathy, glomerulosclerosis, chronic renal failure, diabetic neuropathy, syndrome X, premenstrual syndrome, coronary heart disease, angina pectoris, myocardial infarction, stroke, vascular restenosis, hyperglycemia, hyperinsulinaemia, hyperlipidaemia, hypertryglyceridemia, insulin resistance, impaired glucose metabolism, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose, obesity, diabetic retinopathy, macular degeneration, cataracts, diabetic nephropathy, glomerulosclerosis, diabetic neuropathy, erectile dysfunction, premenstrual syndrome, skin and connective tissue disorders, foot ulcerations and ulcerative colitis, endothelial dysfunction and impaired vascular compliance. Preferably, said combination may be used for the treatment of hypertension, especially ISH, congestive heart failure, endothelial dysfunction, impaired vascular compliance, IGT and type II diabetes mellitus.

A "disease or condition which may be inhibited by the inhibition of AT₁ receptor" as defined in this application comprises, but is not limited to hypertension, congestive heart failure, diabetes, especially type 2 diabetes mellitus, diabetic retinopathy, macular degeneration, diabetic nephropathy, glomerulosclerosis, chronic renal failure, diabetic neuropathy, syndrome X, premenstrual syndrome, coronary heart disease, angina pectoris, myocardial infarction, stroke, vascular restenosis, endothelial dysfunction and the like.

A "disease or condition which may be inhibited by the enhancement of insulin secretion" as defined in this application comprises, but is not limited to hyperglycemia, hyperinsulinaemia, hyperlipidaemia, hypertrygliceridemia, insulin resistance, impaired glucose metabolism, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose, obesity, diabetic retinopathy, macular degeneration, cataracts, diabetic nephropathy, glomerulosclerosis, diabetic neuropathy, erectile dysfunction, premenstrual syndrome, coronary heart disease, hypertension, angina pectoris, myocardial infarction, stroke, vascular restenosis, skin and connective tissue disorders, foot ulcerations and ulcerative colitis, endothelial dysfunction and impaired vascular compliance.

A "disease or condition that may be inhibited by insulin sensitization" as defined in this application comprises, but is not limited to hyperglycemia, hyperinsulinaemia, hyperlipidaemia, hypertrygliceridemia, insulin resistance, impaired glucose metabolism, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose, obesity, diabetic retinopathy, macular degeneration, cataracts, diabetic nephropathy, glomerulosclerosis, diabetic neuropathy, erectile dysfunction, premenstrual syndrome, coronary heart disease, hypertension, angina pectoris, myocardial infarction, stroke, vascular restenosis, skin and connective tissue disorders, foot ulcerations and ulcerative colitis, endothelial dysfunction and impaired vascular compliance.

Hypertension, in connection with a "disease or condition which may be inhibited by the inhibition of AT₁ receptor", a "disease or condition which may be inhibited by the enhancement of insulin secretion", a "disease or condition that may be inhibited by insulin sensitization" includes and is not limited to mild, moderate and severe hypertension as defined in Journal of Hypertension 1999, 17:151-183, especially on page 162. Especially preferred is "isolated systolic hypertension" (ISH).

Preferably, the jointly therapeutically effective amounts of the active agents according to the combination of the present invention can be administered simultaneously or sequentially in any order, e.g. separately or in a fixed combination.

Under certain circumstances, drugs with different mechanisms of action may be combined. However, just considering any combination of drugs having different modes of action but acting in the similar field does not necessarily lead to combinations with advantageous effects.

All the more surprising is the experimental finding that the combined administration of an AT₁ receptor antagonist and insulin secretion enhancer and/or an insulin sensitizer, or, in each case, a pharmaceutically acceptable form thereof, results not only in a beneficial, especially a potentiating or a synergistic, therapeutic effect. Independent thereof, additional benefits resulting from combined treatment can be achieved such as a surprising prolongation of efficacy, a broader variety of therapeutic treatment and surprising beneficial effects on diseases and conditions associated with diabetes, e.g. less gain of weight. An additional and preferred aspect of the present invention is the prevention, delay of progression or treatment of the condition of isolated systolic hypertension and impaired vascular compliance which means decreased vascular elasticity.

The term "potentiation" shall mean an increase of a corresponding pharmacological activity or therapeutical effect, respectively. Potentiation of one component of the combination according to the present invention by co-administration of an other component according to the present invention means that an effect is being achieved that is greater than that achieved with one component alone.

The term "synergistic" shall mean that the drugs, when taken together, produce a total joint effect that is greater than the sum of the effects of each drug when taken alone.

ISH is the most common form of hypertension in people over 50 years. It is defined as elevated systolic blood pressure (above 140 mm Hg) in conjunction with normal diastolic blood pressure (below 90 mm Hg). Elevated systolic blood pressure is an independent risk factor for cardiovascular diseases and may lead e.g. to myocardial hypertrophy and heart failure. ISH is furthermore characterized by an increased pulse pressure, defined as the

difference between systolic and diastolic blood pressures. Elevated pulse pressure is being recognized as the type of hypertension the least likely to be well controlled. A reduction of elevated systolic blood pressure and correspondingly of pulse pressure is associated with a significant risk reduction in cardiovascular death. It has surprisingly been found that the combination of an AT₁ receptor antagonist and an insulin secretion enhancer or an insulin sensitizer leads to a decrease of ISH and pulse rate, both in hypertensive patients having type 2 diabetes mellitus and in hypertensive patient that do not have type 2 diabetes mellitus.

Furthermore, it has been found that the chronic co-administration of either an insulin sensitizer or an insulin secretion enhancer imparts the beneficial effect on blood vessel morphology and function and results in a decrease of vascular stiffness and correspondingly in a maintenance and in an improvement of vascular compliance.

Accordingly, it has been found that the addition of an insulin sensitizer and/or an insulin secretion enhancer to that of an AT₁ receptor antagonist would potentiate the effect on systolic blood pressure and further improve vascular stiffness/compliance. Conversely, the proven antihypertensive effects of an AT₁ receptor antagonist on systolic and diastolic blood pressure may be potentiated by the addition of an insulin sensitizer and/or an insulin secretion enhancer. The benefit of these combinations may also extend to an additional or potentiated effect on endothelial function, and improve vascular function and structure in various organs/tissues including the kidney, heart, eye and brain. Through the reduction in glucose levels, an anti-thrombotic and anti-atherosclerotic effect can also be demonstrated. Reduction of glucose would prevent or minimize the glycosylation of any structural or functional protein within the cardio-renal system. This effect proves to be highly beneficial by evoking an additive or synergistic effect on vascular function/structure when administered with an AT₁ receptor antagonist which alone improves cardiovascular function and structure through a distinct mechanism.

Further benefits are that lower doses of the individual drugs to be combined according to the present invention can be used to reduce the dosage, for example, that the dosages need not only often be smaller but are also applied less frequently, or can be used in order to diminish the incidence of side effects. This is in accordance with the desires and requirements of the patients to be treated.

For example, it has turned out that the combination according to the present invention provides benefit especially in the treatment of modest hypertension or isolated systolic hypertension that is beneficial to all diabetic patients regardless of their hypertensive status, e.g. reducing the risk of negative cardiovascular events by two different modes of action.

The AT₁ receptor antagonists, especially valsartan, have proven to be also useful in the treatment of type 2 diabetes mellitus beyond the reduction of blood pressure in for example improving microalbuminuria. At sub-therapeutic doses, with respect to the treatment of hypertension, the combination according to the invention may be merely used for the treatment of diabetes, especially type 2 diabetes mellitus. In view of the reduced dose of the AT₁ receptor antagonist, there is a considerable safety profile of the combination making it suitable for first line therapy.

The present invention relates to the use of a combination comprising as active ingredients

- (i) an AT₁-receptor antagonist or a pharmaceutically acceptable salt thereof;
- (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof;

for the manufacture of a medicament for the prevention, delay of progression or treatment of a disease and disorder which may be inhibited by the inhibition of AT₁ receptor and by the enhancement of insulin secretion, for example, for the prevention, delay of progression or treatment of hypertension, especially modest hypertension, ISH, congestive heart failure, endothelial dysfunction, impaired vascular compliance, IGT and type II diabetes mellitus.

The present invention also relates to a method for the prevention, delay of progression or treatment of a disease and disorder which may be inhibited by the inhibition of AT₁ receptor and/or by the enhancement of insulin secretion comprising administering to a warm-blooded animal, including man, in need thereof jointly therapeutically effective amounts of

- (i) an AT₁-receptor antagonist or a pharmaceutically acceptable salt thereof;
- (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof.

The pharmaceutical composition according to the present invention as described hereinbefore and hereinafter may be used for simultaneous use or sequential use in any order, for separate use or as a fixed combination.

The pharmaceutical composition according to the present invention comprises a "kit of parts" in the sense that the components can be dosed independently or by use of different fixed combinations with distinguished amounts of the components at different time points. The parts of the "kit of parts" can then e.g. be administered simultaneously or chronologically staggered, that is at different time points and with equal or different time intervals for any part of the "kit of parts". Preferably, the time intervals are chosen such that the effect on the treated disease or condition in the combined use of the parts is larger than the effect that would be obtained by use of only any one of the components. Preferably, there is at least one beneficial effect, e.g. a mutual enhancing of the effect of

- (i) an AT_1 -receptor antagonist or a pharmaceutically acceptable salt thereof;
- (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof;

in particular a potentiation or a synergism, e.g. a more than additive effect, additional advantageous effects, less side effects, a combined therapeutical effect in a non-effective dosage of one or each of the components, especially a potentiation or a strong synergism.

The invention furthermore relates to a commercial package comprising the combination according to the present invention together with instructions for simultaneous, separate or sequential use.

These pharmaceutical preparations are for enteral, such as oral, and also rectal or parenteral, administration to homeotherms, with the preparations comprising the pharmacological active compound either alone or together with customary pharmaceutical auxiliary substances. For example, the pharmaceutical preparations consist of from about 0.1 % to 90 %, preferably of from about 1 % to about 80 %, of the active compound. Pharmaceutical preparations for enteral or parenteral, and also for ocular, administration are, for example, in unit dose forms, such as coated tablets, tablets, capsules or suppositories and also ampoules. These are prepared in a manner that is known per se, for example using conventional mixing, granulation, coating, solubilizing or lyophilizing processes. Thus, pharmaceutical preparations for oral use can be obtained by combining

the active compound with solid excipients, if desired granulating a mixture which has been obtained, and, if required or necessary, processing the mixture or granulate into tablets or coated tablet cores after having added suitable auxiliary substances.

The dosage of the active compound can depend on a variety of factors, such as mode of administration, homeothermic species, age and/or individual condition.

Preferred dosages for the active ingredients of the pharmaceutical combination according to the present invention are therapeutically effective dosages, especially those which are commercially available.

Normally, in the case of oral administration, an approximate daily dose of from about 1 mg to about 360 mg is to be estimated e.g. for a patient of approximately 75 kg in weight.

The dosage of the active compound can depend on a variety of factors, such as mode of administration, homeothermic species, age and/or individual condition.

Valsartan, as a representative of the class of AT₁-receptor antagonists, will be supplied in the form of suitable dosage unit form, for example, a capsule or tablet, and comprising a therapeutically effective amount, e.g. from about 20 to about 320 mg, of valsartan which may be applied to patients. The application of the active ingredient may occur up to three times a day, starting e.g. with a daily dose of 20 mg or 40 mg of valsartan, increasing via 80 mg daily and further to 160 mg daily up to 320 mg daily. Preferably, valsartan is applied twice a day with a dose of 80 mg or 160 mg, respectively, each. In a low-doses formulation, valsartan with a dose of 20 mg or 40 mg may be used. Corresponding doses may be taken, for example, in the morning, at mid-day or in the evening. Preferred is b.i.d. administration.

The insulin secretion enhancer nateglinide (I) is preferably administered to the warm-blooded animal in a dosage in the range of about 5 to 1200, more preferably 25 to 800, mg/day, when the warm-blooded animal is a human of about 70 kg body weight. Preferred dosages contain 30mg, 60mg, 120 mg or 180 mg of nateglinide to be administered preferably before the main meals. In a low dose combination, the dosage of nateglinide to be administered preferably is 30 mg, 40 mg or furthermore 60 mg. Depending on the

number of main meals the dose regimen are two times a day (BID) or three times a day (TID) or four times a day (QID).

The insulin secretion enhancer repaglinde is preferably administered in a dosage range of about 0.01 mg to about 8 mg, more preferred from about 0.5 to about 6 mg.

The insulin sensitizer metformin is preferably administered in a dosage range of about 100 mg to about 1200 mg per dose unit, especially 500 mg, 850 mg or 1000 mg. In a low dose combination, metformin is preferably administered in a dosage of 125 mg, 250 mg or 500 mg.

Formulation Example 1:

Film-Coated Tablets:

Components	Composition Per Unit (mg)	Standards
Granulation		
Valsartan [= active ingredient]	80.00	
Microcrystalline cellulose/ Avicel PH 102	54.00	NF, Ph. Eur
Crospovidone	20.00	NF, Ph. Eur
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	0.75	Ph. Eur/ NF
Magnesium stearate	2.5	NF, Ph. Eur
Blending		
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	0.75	Ph. Eur/ NF
Magnesium stearate	2.00	NF, Ph. Eur
Coating		
Purified water ¹⁾	-	
DIOLACK pale red 00F34899	7.00	
Total tablet mass	167.00	

¹⁾ Removed during processing.

The film-coated tablet is manufactured e.g. as follows:

A mixture of valsartan, microcrystalline cellulose, crospovidone, part of the colloidal anhydrous silica/colloidal silicon dioxide/Aerosil 200, silicon dioxide and magnesium stearate is premixed in a diffusion mixer and then sieve through a screening mill. The resulting mixture is again pre-mixed in a diffusion mixer, compacted in a roller compacter and then sieve through a screening mill. To the resulting mixture, the rest of the colloidal anhydrous silica/colloidal silicon dioxide/Aerosil 200 are added and the final blend is made

in a diffusion mixer. The whole mixture is compressed in a rotary tableting machine and the tablets are coated with a film by using Diolack pale red in a perforated pan.

Formulation Example 2:

Film-coated tablets:

Components	Composition Per Unit (mg)	Standards
Granulation		
Valsartan [= active ingredient]	160.00	
Microcrystalline cellulose/ Avicel PH 102	108.00	NF, Ph. Eur
Crospovidone	40.00	NF, Ph. Eur
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	1.50	Ph. Eur/ NF
Magnesium stearate	5.00	NF, Ph. Eur
Blending		
Colloidal anhydrous silica / colloidal silicon dioxide / Aerosil 200	1.50	Ph. Eur/ NF
Magnesium stearate	4.00	NF, Ph. Eur
Coating		
Opadry Light Brown 00F33172	10.00	
Total tablet mass	330.00	

The film-coated tablet is manufactured e.g. as described in Formulation Example 1.

Formulation Example 3:

Film-Coated Tablets:

Components	Composition Per Unit (mg)	Standards
Core: Internal phase		
Valsartan [= active ingredient]	40.00	
Silica, colloidal anhydrous (Colloidal silicon dioxide) [= Glidant]	1.00	Ph. Eur, USP/NF
Magnesium stearate [= Lubricant]	2.00	USP/NF
Crospovidone [Disintegrant]	20.00	Ph. Eur
Microcrystalline cellulose [= Binding agent]	124.00	USP/NF
External phase		
Silica, colloidal anhydrous, (Colloidal silicon dioxide) [= Glidant]	1.00	Ph. Eur, USP/NF
Magnesium stearate [Lubricant]	2.00	USP/NF
Film coating		
Opadry® brown OOF 16711 ¹⁾	9.40	
Purified Water ²⁾	-	
Total tablet mass	199.44	

¹⁾ The composition of the Opadry® brown OOF16711 coloring agent is tabulated below.

²⁾ Removed during processing

Opadry® Composition:

Ingredient	Approximate % Composition
Iron oxide, black (C.I. No. 77499, E 172)	0.50
Iron oxide, brown (C.I. No. 77499, E 172)	0.50
Iron oxide, red (C.I. No. 77491, E 172)	0.50
Iron oxide, yellow (C.I. No. 77492, E 172)	0.50
Macrogolum (Ph. Eur)	4.00
Titanium dioxide (C.I. No. 77891, E 171)	14.00
Hypromellose (Ph. Eur)	80.00

The film-coated tablet is manufactured e.g. as described in Formulation Example 1.

Formulation Example 4:

Capsules:

Components	Composition Per Unit (mg)
Valsartan [= active ingredient]	80.00
Microcrystalline cellulose	25.10
Crospovidone	13.00
Povidone	12.50
Magnesium stearate	1.30
Sodium lauryl sulphate	0.60
Shell	
Iron oxide, red (C.I. No. 77491, EC No. E 172)	0.123
Iron oxide, yellow (C.I. No. 77492, EC No. E 172)	0.123
Iron oxide, black (C.I. No. 77499, EC No. E 172)	0.245
Titanium dioxide	1.540
Gelatin	74.969
Total tablet mass	209.50

The tablet is manufactured e.g. as follows:

Granulation/Drying

Valsartan and microcrystalline cellulose are spray-granulated in a fluidised bed granulator with a granulating solution consisting of povidone and sodium lauryl sulphate dissolved in purified water. The granulate obtained is dried in a fluidised bed dryer.

Milling/Blending

The dried granulate is milled together with crospovidone and magnesium stearate. The mass is then blended in a conical screw type mixer for approximately 10 minutes.

Encapsulation

The empty hard gelatin capsules are filled with the blended bulk granules under controlled temperature and humidity conditions. The filled capsules are dedusted, visually inspected, weightchecked and quarantined until by Quality assurance department.

Formulation Example 5:

Capsules:

Components	Composition Per Unit (mg)
Valsartan [= active ingredient]	160.00
Microcrystalline cellulose	50.20
Crospovidone	26.00
Povidone	25.00
Magnesium stearate	2.60
Sodium lauryl sulphate	1.20
Shell	
Iron oxide, red (C.I. No. 77491, EC No. E 172)	0.123
Iron oxide, yellow (C.I. No. 77492, EC No. E 172)	0.123
Iron oxide, black (C.I. No. 77499, EC No. E 172)	0.245
Titanium dioxide	1.540
Gelatin	74.969
Total tablet mass	342.00

The formulation is manufactured e.g. as described in Formulation Example 4.

Formulation Example 6:

Hard Gelatine Capsule:

Components	Composition Per Unit (mg)
Valsartan [= active ingredient]	80.00
Sodium laurylsulphate	0.60
Magnesium stearate	1.30
Povidone	12.50
Crospovidone	13.00
Microcrystalline cellulose	21.10
Total tablet mass	130.00

Examples 7 to 11:

Example	7	8	9	10	11
Components	Composition per unit (mg)	Composit ion per unit (mg)	Composit ion per unit (mg)	Composit ion per unit (mg)	Composit ion per unit (mg)
Granulation					
Valsartan Drug Substance (DS)	80.000	160.000	40.000	320.000	320.000
Microcrystalline Cellulose (NF, Ph.Eur.)/ Avicel PH 102	54.000	108.000	27.000	216.000	216.000
Croscopovidone (NF, Ph.Eur.)	15.000	30.000	7.500	80.000	60.000
Colloidal Anhydrous Silica (Ph. Eur.)/Colloidal Silicon Dioxide (NF)/Aerosil 200	1.500	3.000	0.750	3.000	6.000
Magnesium Stearate (NF, Ph.Eur.)	3.000	6.000	1.500	10.000	12.000
Blending					
Colloidal Anhydrous Silica (Ph. Eur.)/Colloidal Silicon Dioxide (NF)/Aerosil 200	---	---	---	3.000	-
Magnesium Stearate, NF, Ph.Eur.	1.500	3.000	0.750	8.000	6.000
Core Weight/mg	155.000	310.000	77.500	640.000	620.000
Coating	-	-	3.800	15.000	16.000

Example 12 :

108,000 tablets, each which contain 120 mg of nateglinide are prepared as follows:

<u>Composition:</u>	nateglinide	12.960 kg
	lactose, NF	30.564 kg
	microcrystalline cellulose, NF	15.336 kg
	povidone, USP	2.592 kg
	croscarmellose sodium, NF	3.974 kg
	colloidal silicon dioxide, NF	1.382 kg
	magnesium stearate, NF	1.231 kg
	coating: opadry yellow	1.944 kg
	purified water, USP*	Q.S.

*: removed during process

Preparation process: The microcrystalline cellulose, povidone, part of the croscarmellose sodium, nateglinide and lactose are mixed in a high shear mixer and afterwards granulated using purified water. Alternatively, the microcrystalline cellulose, povidone, a portion of the croscarmellose sodium, nateglinide and lactose are granulated in a collette gral granulator with the addition of purified water. The wet granules are dried in a fluid bed dryer and passed through a screen. The colloidal silicon dioxide and the rest of the croscarmellose sodium are mixed, passed through a screen and blended with the dried granules in a V-blender. The magnesium stearate is passed through a screen, blended with the blend from the V-blender and afterwards the total mixture is compressed to tablets. The opadry yellow is suspended in purified water and the tablets are coated with the coating suspension.

Examples 13-15:

Component	60mg	120mg	180mg
Starlix DS (H-form crystal modification)	60	120	180
Lactose Monohydrate	141.5	283	214
Microcrystalline Cellulose	71	142	107
Povidone K30	12	24	23
Croscarmellose Sodium	12	24	34
Sub-Total (Granulation)	296.5	593	558
Croscarmellose Sodium	6.4	12.8	24.5
Colloidal Silicone Dioxide	6.4	12.8	12.3
Magnesium Stearate	5.7	11.4	15.2
Sub-Total (Core)	(315)	(630)	(610)
Opadry	9	18	18
Total	324	648	628

What is claimed is

1. A pharmaceutical composition, comprising as active ingredients
 - (i) an AT₁-receptor antagonist or a pharmaceutically acceptable salt thereof;
 - (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof; and
 - (iii) a pharmaceutically acceptable carrier.
2. A composition according to claim 1 wherein the AT₁ receptor antagonist is selected from the group consisting of valsartan, losartan, candesartan, eprosartan, irbesartan, olmesartan, tasosartan, and telmisartan, and, in each case, a pharmaceutically acceptable salt thereof.
3. A composition according to claim 2 wherein the AT₁ receptor antagonist is valsartan or a pharmaceutically acceptable salt thereof.
4. A composition according to claim 1 wherein the insulin secretion enhancer is selected from tolbutamide; chlorpropamide; tolazamide; acetohexamide; glycopyramide; glibenclamide; gliclazide; 1-butyl-3-metanilylurea; carbutamide; glibonuride; glipizide; gliquidone; glisoxepid; glybuthiazole; glibuzole; glyhexamide; glymidine; glypinamide; phenbutamide; tolylcyclamide, nateglinide repaglinide; mitiglinide; glimepiride; and, in each case, a pharmaceutically acceptable salt thereof.
5. A composition according to claim 4 wherein the insulin secretion enhancer is nateglinide or a pharmaceutically acceptable salt thereof.
6. A composition according to claim 1 wherein the insulin sensitizer is metformin or a pharmaceutically acceptable salt thereof.
7. A composition according to any one of claims 1 to 6 for the prevention, delay of progression or treatment of a of disease and disorder selected from the group consisting of hypertension, congestive heart failure, diabetes, especially type 2 diabetes mellitus, diabetic retinopathy, macular degeneration, diabetic nephropathy, glomerulosclerosis, chronic renal

failure, diabetic neuropathy, syndrome X, premenstrual syndrome, coronary heart disease, angina pectoris, myocardial infarction, stroke, vascular restenosis, hyperglycemia, hyperinsulinaemia, hyperlipidaemia, hypertriglyceridemia, insulin resistance, impaired glucose metabolism, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose, obesity, diabetic retinopathy, macular degeneration, cataracts, diabetic nephropathy, glomerulosclerosis, diabetic neuropathy, erectile dysfunction, skin and connective tissue disorders, foot ulcerations and ulcerative colitis, endothelial dysfunction and impaired vascular compliance.

8. A composition according to any one of claims 1 to 7 for the prevention, delay of progression or treatment of a of disease and disorder selected from the group consisting of hypertension, especially ISH, congestive heart failure, endothelial dysfunction, impaired vascular compliance, IGT and type II diabetes mellitus.

9. A method for the prevention, delay of progression or treatment of a disease and disorder which may be inhibited by the inhibition of AT_1 receptor and/or by the enhancement of insulin secretion comprising administering to a warm-blooded animal, including man, in need thereof jointly therapeutically effective amounts of

- (i) an AT_1 -receptor antagonist or a pharmaceutically acceptable salt thereof;
- (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof.

10. Use of a combination comprising as active ingredients

- (i) an AT_1 -receptor antagonist or a pharmaceutically acceptable salt thereof;
- (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or
(b) an insulin sensitizer or a pharmaceutically acceptable salt thereof;

for the manufacture of a medicament for the prevention, delay of progression or treatment of a disease and disorder which may be inhibited by the inhibition of AT_1 receptor and by the enhancement of insulin secretion.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
28 February 2002 (28.02.2002)

PCT

(10) International Publication Number
WO 02/015933 A3

- (51) International Patent Classification⁷: **A61K 31/41**,
A61P 5/00 // (A61K 31/41, 31:64) (A61K 31/41, 31:195)
(A61K 31/41, 31:155)
- (21) International Application Number: PCT/EP01/09587
- (22) International Filing Date: 20 August 2001 (20.08.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/643,641 22 August 2000 (22.08.2000) US
- (71) Applicant (*for all designated States except AT, US*): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).
- (71) Applicant (*for AT only*): **NOVARTIS PHARMA GMBH** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **ALLISON, Malcolm** [GB/CH]; Grenzacherstrasse 108, CH-4058 Basel (CH). **GATLIN, Marjorie, Regan** [US/US]; 913 Lovering Avenue, Wilmington, DE 19806 (US).
- (74) Agent: **BECKER, Konrad**; Novartis AG, Corporate Intellectual Property, Patent & Trademark Dept., CH-4002 Basel (CH).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (88) Date of publication of the international search report:
14 August 2003
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/015933 A3

(54) Title: COMBINATION COMPRISING AN AT₁-RECEPTOR ANTAGONIST AND AN INSULIN SECRETION ENHANCER OR AN INSULIN SENSITISER

(57) Abstract: The present invention relates to a combination, especially a pharmaceutical composition, comprising as active ingredients (i) an AT₁-receptor antagonist or a pharmaceutically acceptable salt thereof; (ii) (a) an insulin secretion enhancer or a pharmaceutically acceptable salt thereof or (b) an insulin sensitizer or a pharmaceutically acceptable salt thereof; and, in case of a pharmaceutical composition, a pharmaceutically acceptable carrier.

INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/EP 01/09587

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/41 A61P5/00 //(A61K31/41,31:64), (A61K31/41,31:195),
(A61K31/41,31:155)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data, EMBASE, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 45818 A (CAMERON NORMAN EUGENE ;UNIV ABERDEEN (GB); ASTRAZENECA UK LTD (GB)) 10 August 2000 (2000-08-10) claims 7,8,10,13 ---	1-4,7-10
P,X	WO 01 21602 A (DEVASTHALE PRATIK ;SQUIBB BRISTOL MYERS CO (US); CHEN SEAN (US); J) 29 March 2001 (2001-03-29) page 16; claims 40,48 -----	1-5,7-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

9 December 2002

Date of mailing of the international search report

10. 04 2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

INTERNATIONAL SEARCH REPORT

international application No.
PCT/EP 01/09587

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 9 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: partially 1-5,7-10
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

4, 5 completely, 1-3, 7-10 partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 4,5 complete, 1-3,7-10 partially

Composition comprising an AT1 (angiotensin II) receptor antagonist and an insulin secretion enhancer for the treatment of diabetes, obesity, hypertension.

2. Claims: 6 complete, 1-3,7-10 partially

Composition comprising an AT1 (angiotensin II) receptor antagonist and an insulin sensitizer for the treatment of diabetes, obesity, hypertension.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: partially 1-5,7-10

Present claims 1-5,7-10 relate to compounds/compositions defined by reference to desirable characteristics or properties, namely: "AT1 receptor antagonist", "insulin secretion enhancer", "insulin sensitizer" (claims 1,9,10)

The claims cover all compounds/compositions having these characteristics or properties, whereas the application provides support within the meaning of Article 6 PCT and disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compounds by reference to a result to be achieved: "AT1 receptor antagonist", "insulin secretion enhancer", "insulin sensitizer". Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

Consequently, the search for the first invention has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds specifically mentioned by chemical name in claims 2-5 with due regard to the general idea underlying the present application.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No

PCT/EP 01/09587

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 0045818	A	10-08-2000	AU	2304700 A		25-08-2000
			BR	0007996 A		30-10-2001
			CN	1338937 T		06-03-2002
			CZ	20012806 A3		13-02-2002
			EE	200100405 A		15-10-2002
			EP	1150678 A1		07-11-2001
			WO	0045818 A1		10-08-2000
			HU	0105138 A2		29-04-2002
			JP	2002536332 T		29-10-2002
			NO	20013812 A		02-10-2001
			PL	350312 A1		02-12-2002
			SK	11102001 A3		09-05-2002
			TR	200102229 T2		21-12-2001

WO 0121602	A	29-03-2001	AU	7593500 A		24-04-2001
			BR	0014189 A		20-08-2002
			CN	1374955 T		16-10-2002
			EP	1218361 A1		03-07-2002
			JP	2003509503 T		11-03-2003
			NO	20021408 A		14-05-2002
			TR	200200732 T2		21-10-2002
			WO	0121602 A1		29-03-2001
			US	6414002 B1		02-07-2002

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.